


Domestic Violence

1 in 3 women

throughout the world will experience physical and/or sexual violence by a partner or sexual violence by a non-partner



◆ Key Facts:

- ✚ Violence against women - particularly intimate partner violence and sexual violence - are major public health problems and violations of women's human rights.
- ✚ Recent global prevalence figures indicate that about 1 in 3 (35%) of women worldwide have experienced either physical and/or sexual intimate partner violence or non-partner sexual violence in their lifetime.
- ✚ Globally, as many as 38% of murders of women are committed by an intimate partner.
- ✚ Violence can negatively affect women's physical, mental, sexual and reproductive health, and may increase vulnerability to HIV.
- ✚ Factors associated with increased risk of perpetration of violence include low education, child maltreatment or exposure to violence in the family, **harmful use of alcohol**, attitudes accepting of violence and gender inequality.
- ✚ There is evidence from high-income settings that school-based programs may be effective in preventing relationship violence (or dating violence) among young people.
- ✚ In low-income settings, primary prevention strategies, such as microfinance combined with gender equality training and community-based initiatives that address gender inequality and relationship skills, hold promise.

Domestic Violence

◆ Introduction

The United Nations defines violence against women as "any act of gender-based violence that results in, or is likely to result in, physical, sexual or mental harm or suffering to women, including threats of such acts, coercion or arbitrary deprivation of liberty, whether occurring in public or in private life."

Intimate partner violence refers to behavior by an intimate partner or ex-partner that causes physical, sexual or psychological harm, including physical aggression, sexual coercion, and psychological abuse and controlling behaviors.

Sexual violence is "any sexual act, attempt to obtain a sexual act, or other act directed against a person's sexuality using coercion, by any person regardless of their relationship to the victim, in any setting. It includes rape, defined as the physically forced or otherwise coerced penetration of the vulva or anus with a penis, other body part or object."

BOX 1: **Extent of intimate partner violence**

Most reported intimate partner violence is perpetrated by men towards women (1). However, violence is also committed by women towards men (2) and within same sex relationships (3). Variations in methodologies and definitions of violence between surveys make the extent of intimate partner violence and differences between countries hard to estimate. However, the *WHO multi-country study on women's health and domestic violence against women* (4), one of the few

studies to report comparable data, shows that between 15% (Japan) and 71% (Ethiopia) of women reported experiencing physical and sexual violence by an intimate partner over their lifetime, and between 3.8% (Japan) and 53% (Ethiopia) as experiencing such violence within the past year. In a survey of 24,000 men and women in Canada, 7% of women and 6% of men reported having been victims of intimate partner violence in the last five years (5).

◆ Scope of the Problem

Population-level surveys based on reports from victims provide the most accurate estimates of the prevalence of intimate partner violence and sexual violence in non-conflict settings. The first report of the "WHO Multi-country study on women's health and domestic violence against women" (2005) in 10 mainly low- and middle-income countries found that, among women aged 15-49:

Domestic Violence

- ✚ between 15% of women in Japan and 71% of women in Ethiopia reported physical and/or sexual violence by an intimate partner in their lifetime;
- ✚ between 0.3–11.5% of women reported sexual violence by someone other than a partner since the age of 15 years;
- ✚ the first sexual experience for many women was reported as forced – 17% of women in rural Tanzania, 24% in rural Peru, and 30% in rural Bangladesh reported that their first sexual experience was forced.

A more recent analysis of WHO with the London School of Hygiene and Tropical Medicine and the Medical Research Council, based on existing data from over 80 countries, found that globally 35% of women have experienced either physical and/or sexual intimate partner violence or non-partner sexual violence. Most of this violence is intimate partner violence. Worldwide, almost one-third (30%) of all women who have been in a relationship have experienced physical and/or sexual violence by their intimate partner, in some regions this is much higher. Furthermore, globally as many as 38% of all murders of women are committed by intimate partners.

Intimate partner and sexual violence are mostly perpetrated by men against women. Child sexual abuse affects both boys and girls. International studies reveal that approximately 20% of women and 5–10% of men report being victims of sexual violence as children. Violence among young people, including dating violence, is also a major problem.

◆ Risk Factors

Factors associated with intimate partner and sexual violence occur at individual, family, community, and wider society levels. Some factors are associated with being a perpetrator of violence, some are associated with experiencing violence, and some are associated with both.

1. Risk factors for both intimate partner and sexual violence include:

- ✚ lower levels of education (perpetration of sexual violence and experience of sexual violence);
- ✚ exposure to child maltreatment (perpetration and experience);
- ✚ witnessing family violence (perpetration and experience);
- ✚ antisocial personality disorder (perpetration);
- ✚ harmful use of alcohol (perpetration and experience);
- ✚ having multiple partners or suspected by their partners of infidelity (perpetration); and
- ✚ attitudes that are accepting of violence and gender inequality (perpetration and experience).

Domestic Violence

2. Factors specifically associated with intimate partner violence include:

- ✚ past history of violence;
- ✚ marital discord and dissatisfaction;
- ✚ difficulties in communicating between partners.

3. Factors specifically associated with sexual violence perpetration include:

- ✚ beliefs in family honor and sexual purity
- ✚ ideologies of male sexual entitlement and
- ✚ weak legal sanctions for sexual violence.

◆ Health Consequences

Intimate partner and sexual violence have serious short- and long-term physical, mental, sexual and reproductive health problems for survivors and for their children, and lead to high social and economic costs.

- ✚ Violence against women can have fatal results like homicide or suicide.
- ✚ It can lead to injuries, with 42% of women who experience intimate partner violence reporting an injury as a consequence of this violence.
- ✚ Intimate partner violence and sexual violence can lead to unintended pregnancies, induced abortions, gynecological problems, and sexually transmitted infections, including HIV. The 2013 analysis found that women who had been physically or sexually abused were 1.5 times more likely to have a sexually transmitted infection and, in some regions, HIV, compared to women who had not experienced partner violence. They are also twice as likely to have an abortion.
- ✚ Intimate partner violence in pregnancy also increases the likelihood of miscarriage, stillbirth, pre-term delivery and low birth weight babies.
- ✚ These forms of violence can lead to depression, post-traumatic stress disorder, sleep difficulties, eating disorders, emotional distress, and suicide attempts. The same study found that women who have experienced intimate partner violence were almost twice as likely to experience depression and problem drinking. The rate was even higher for women who had experienced non-partner sexual violence.
- ✚ Health effects can also include headaches, back pain, abdominal pain, fibromyalgia, gastrointestinal disorders, limited mobility, and poor overall health.
- ✚ Sexual violence, particularly during childhood, can lead to increased smoking, drug and alcohol misuse, and risky sexual behaviors in later life. It is also associated with perpetration of violence (for males) and being a victim of violence (for females).

Domestic Violence

◆ Impact on Children

Children who grow up in families where there is violence may suffer a range of behavioral and emotional disturbances. These can also be associated with perpetrating or experiencing violence later in life. Intimate partner violence has also been associated with higher rates of infant and child mortality and morbidity (e.g. diarrheal disease, malnutrition).

◆ Social and Economic Costs

The social and economic costs of intimate partner and sexual violence are enormous and have ripple effects throughout society. Women may suffer isolation, inability to work, loss of wages, lack of participation in regular activities, and limited ability to care for themselves and their children.

◆ Prevention and Response

Currently, there are few interventions whose effectiveness have been proven through well- designed studies. More resources are needed to strengthen the prevention of intimate partner and sexual violence, including primary prevention, i.e. stopping it from happening in the first place.

Regarding primary prevention, there is some evidence from high-income countries that school-based programs to prevent violence within dating relationships have shown effectiveness. However, these have yet to be assessed for use in resource-poor settings. Several other primary prevention strategies: those that combine microfinance with gender equality training, that promote communication and relationship skills within couples and communities, that reduce access to, and harmful use of alcohol, and that change cultural gender norms, have shown some promise but need to be evaluated further.

To achieve lasting change, it is important to enact legislation and develop policies that:

- ✚ address discrimination against women;
- ✚ promote gender equality;
- ✚ support women; and
- ✚ help to move towards more peaceful cultural norms.

An appropriate response from the health sector can play an important role in the prevention of violence. Sensitization and education of health and other service providers is therefore another important strategy. To address fully the consequences of violence and the needs of victims/survivors requires a multi-sectoral response.

Source: http://www.who.int/topics/gender_based_violence/en/